



1  
3  
5  
7  
9 **Research Article**

11 **Immunocontraception and Increased**  
13 **Longevity in Equids**

15 **J.F. Kirkpatrick,<sup>1\*</sup> and A. Turner<sup>2</sup>**

17 <sup>1</sup>*The Science and Conservation Center, Billings, Montana*

19 <sup>2</sup>*Assateague Island National Seashore, Berlin, Maryland*

21 Intensive population management by means of fertility control has been shown  
23 to change the age profile of a wild horse herd. The primary change has been  
25 an increase in the number and percent of older animals, as expected, but also the  
27 appearance of new and older age classes. An examination of direct effects of  
29 fertility control on two groups of treated animals shows a significant increase in  
31 longevity over non-treated animals that is associated with contraceptive  
treatment. The mean age at death (MAD) was calculated for 128 wild horses  
for which precise birth and death dates were known, including 56 stallions,  
42 untreated mares, 11 mares treated with a porcine zona pellucida contraceptive  
vaccine for 1-2 years, and 19 mares treated with the same vaccine for  $\geq 3$  years.  
The MAD for stallions ( $10.3 \pm 0.84$  [SEM] years), and mares treated for 1-2 years  
( $10.2 \pm 0.56$ ), was significantly greater ( $P < 0.05$ ) than for untreated mares  
( $6.4 \pm 0.85$ ), and significantly  $< 19.9 \pm 1.66$  for mares treated  $\geq 3$  years  
( $19.9 \pm 1.66$ ). Zoo Biol 25:1-8, 2007. © 2006 Wiley-Liss, Inc.

33 **Keywords:** contraception; equids; horses; immunocontraception; longevity; horses

35 **INTRODUCTION**

37 The application of porcine zona pellucida (PZP) immunocontraception to  
39 captive and free-ranging wildlife, for population management has increased since its  
initial use for this purpose in 1988 [Kirkpatrick et al., 1990]. Previous research has

41 Grant sponsor: National Park Service; Grant number: CA-1600-30003; Grant sponsor: National Institute  
of Health; Grant number: 1 R15 HDZ6898-01A1; Grant sponsor: Assateague Island National Seashore;  
43 Grant sponsor: Science and Conservation Center at ZooMontana.

45 \*Correspondence to: Jay F. Kirkpatrick, The Science and Conservation Center, 2100 South Shiloh Road,  
Billings, MT 59106. E-mail: Jkirkpatrick@montana.net

47 Received 23 January 2006; Accepted 16 May 2006

DOI 10.1002/zoo.20109

Published online in Wiley InterScience (www.interscience.wiley.com).

1 shown the PZP vaccine to be efficacious at both the level of the individual animal  
2 of many species [Kirkpatrick et al., 1990, 1995, 1996; Turner et al., 1996; Shideler  
3 et al., 2002; Frank et al., 2005] and in managing entire populations [Turner and  
4 Kirkpatrick, 2002; Naugle et al., 2002; Deigert et al., 2003; Rutberg et al., 2004] of  
5 captive exotic and native free-ranging species. Additionally, other studies have  
6 documented the vaccine's safety with regard to pregnant animals, the reversibility of  
7 contraceptive action in equids [Kirkpatrick et al., 1991a, 1992, 1995; Kirkpatrick and  
8 Turner, 2002], cervids [Turner et al., 1996; Naugle et al., 2002; Rutberg et al., 2004],  
9 African elephants [Fayrer-Hosken et al., 2000] and the lack of effect on foal survival  
10 or out-of-season births in equids [Kirkpatrick and Turner, 2003].

11 The ability to manage population growth in some manner, either by slowing  
12 growth rates, or achieving zero population growth or even through herd reduction  
13 has been shown after long-term studies [Kirkpatrick and Turner, 2002; Turner and  
14 Kirkpatrick, 2002], but several unanticipated phenomena have modified expectations  
15 for managing populations. First, it was not clear at the outset of research with this  
16 vaccine, 17 years ago, just what long-term health effects the application of the  
17 vaccine might have for individual animals and how these effects might modify  
18 population management. Two of the more significant changes that have been noted  
19 include a significant improvement in body condition among chronically treated wild  
20 mares and an increase in new and older age classes at the population level [Turner  
21 and Kirkpatrick, 2002]. It has been hypothesized that these two phenomena were  
22 related [Turner and Kirkpatrick, 2002].

23 A comparison of age classes in a PZP-treated population of wild horses on  
24 Assateague Island National Seashore (ASIS), in 1990, 1995, and 2001 [Turner and  
25 Kirkpatrick, 2002] showed a steady shift to older age classes in this population. The  
26 causes for this increase in older animals were not surprising and were assumed to be  
27 the result of fewer pregnancies among treated animals and a resulting increase in  
28 body condition, less mortality, and therefore, longer life. The age class shift,  
29 however, was apparent only at the population level and not at the level of individual  
30 animals.

31 It was the purpose of this retrospective study to examine the relationship  
32 between PZP treatment in two groups of animals of differing treatment protocols  
33 and untreated animals, and increased longevity. The working hypothesis was that  
34 increasing levels of contraception were associated with increased longevity in the two  
35 animal cohorts of the study population, and that both treatment groups would show  
36 increased longevity over non-treated animals.

## 37

## 39 MATERIALS AND METHODS

### 41 Study Population

42 The study population was made up of free-ranging wild horses inhabiting  
43 Assateague Island National Seashore in Maryland. This population has inhabited  
44 the island for approximately 330 years at varying densities and population numbers.  
45 In 1965, the Maryland portion of the island, comprising 16,083 hectares was  
46 designated as Assateague Island National Seashore, under the management of the  
47 National Park Service (NPS) and at that time the 28 wild horses inhabiting the  
48 national Seashore were declared a cultural and historic resource and included among



1 the fauna of the island. In 1988, when research with the PZP vaccine was initiated  
2 with 26 mares [Kirkpatrick et al., 1990], the herd population was 129. From 1988  
3 through 1993, 46 different mares were inoculated for varying periods of time as part  
4 of the original research effort, and during that time, the herd size grew to 156.  
5 Beginning in 1994, with the herd size at 166, every remaining untreated mare that  
6 was 2 years or older was given a single "primer" dose of PZP and in 1995, with the  
7 herd size at 173, a contraceptive management plan was developed and initiated, with  
8 an immediate goal of reducing the population to 150 animals. For the first several  
9 years the management plan called for the inoculation of all 2-year old mares, and  
10 booster inoculations of PZP at age 3 and 4, followed by withdrawal of treatment  
11 until the mare produced three living offspring or a second generation, after which the  
12 mare would be treated with the PZP annually until her death. This provided for a  
13 delay in the reproductive age of mares, but a continuous contribution of genes from  
14 all mares on the island and this approach resulted in zero population growth almost  
15 immediately [Turner and Kirkpatrick, 2002]. This plan was modified in 1998 to allow  
16 only two foals after withdrawal from treatment, and again in 2000 to allow only a  
17 single foal, largely because there was no apparent decrease in the population as a  
18 result of contraception. By March 2006, the population had decreased to 140 (a 19%  
19 decrease) without the need for removal of animals.

#### 21 **Vaccine Application**

22 The vaccine was prepared according to the modified method of Dunbar et al.  
23 [1980] at the University of California-Davis from 1988–1997 and at the Science and  
24 Conservation Center thereafter. Delivery of the vaccine was by 1.0 cc Pneu-Darts,  
25 (Pneu-Dart, Inc., Williamsport, PA) at doses of 100 µg, with an appropriate  
26 adjuvant, as described by Kirkpatrick et al. (1990, 1991) and authorized under the  
27 Food and Drug Administration Investigational New Animal Drug Exemption File  
28 8857-G002. No horses were captured or handled during the course of this study.

#### 29 **Pregnancy Testing**

30 To assess pregnancy loss or undetected neonatal losses, 346 horses were  
31 pregnancy-tested by means of urinary or fecal steroid conjugates [Kirkpatrick et al.,  
32 1990, 1991b] between 1988–2005. Fall pregnancy rates based on estrone conjugate  
33 analysis were compared to foals counted the following spring and summer.

#### 35 **Data Collection**

36 All animals in the Assateague population are monitored every other month by  
37 an ASIS employee (AT) and data collected including: 1) identification of individual  
38 animals by unique markings, 2) band affiliation, 3) band location, 4) band habitat  
39 use, 5) the presence and date of new foals, and 6) the date and death of older  
40 animals. Not all carcasses of dead animals are discovered immediately, but the  
41 continued absence of animals from their bands is noted on a monthly basis, thus the  
42 date of death on a yearly basis is standard and in most cases, the date of death can be  
43 noted by month.

44 The data presented in this retrospective study come from records for 128 horses  
45 for whose dates of birth and death are certain, covering the period from 1976–2005.  
46 Only horses that lived to be a minimum of 2 years old were included, to normalize  
47 data to the treated population; on ASIS, PZP treatment is always initially

#### 4 Kirkpatrick and Turner

1 administered at 2 years of age and contraceptive results would not become apparent  
2 until animals were 3 years old. Horses dying from car collisions, accidental death by  
3 deer hunters, and drowning by a tidal surge in 1992 were not included in this study,  
4 thus, only those horses that died from causes unrelated to human activities and  
5 natural disasters were included. The horses were grouped into four cohorts,  
6 including: 1) 56 stallions, 2) 42 mares that had never been treated with PZP, 3) 11  
7 mares that had been treated for up to 2 years, and 4) 19 mares treated for 3 or more  
8 years. The mean age at death (MAD) was calculated for each group and compared  
9 to the MAD of the groups and the differences tested for significance by paired *t*-test.

#### 11 RESULTS

13 The results are presented in Table 1. In general, there was increasing longevity  
14 among individual animals from untreated mares dying at the youngest ages  
15 ( $6.4 \pm 0.85$  years), to stallions, and mares treated up to 2 years dying at intermediate  
16 ages ( $10.3 \pm 0.84$  and  $10.2 \pm 0.56$  years, respectively, to mares treated for  $\geq 3$  years  
17 ( $19.9 \pm 1.66$  years). The mean age at death (MAD) of the 56 stallions was not  
18 significantly different from the MAD of mares treated for  $< 3$  years ( $P = 0.973$ ;  
19  $t = 0.032$ ), but was significantly greater than that of untreated mares ( $P = 0.0006$ ;  
20  $t = 3.53$ ), and significantly less than the MAD of mares treated for  $\geq 3$  years  
21 ( $P = 0.005$ ;  $t = 5.51$ ). The MAD of untreated mares was significantly less than that of  
22 mares treated for  $< 3$  years ( $P = 0.064$ ;  $t = 2.84$ ), and significantly less than the MAD  
23 of mares treated for  $\geq 3$  years ( $P = 0.0001$ ;  $t = 9.67$ ). The MAD of mares treated for  
24  $< 3$  years was significantly less than the MAD of mares treated for  $\geq 3$  years  
25 ( $P = 0.005$ ;  $t = 3.92$ ).

#### 27 DISCUSSION

29 Age-specific reproductive success for the ASIS horses indicates an increasing  
30 trend from age 3 through  $> 6$  years [Keiper and Houpt, 1984]. This pattern is similar  
31 to that seen in other wild horse herds in which reproductive success is low for 1- and  
32 2-year-olds, increasing from 3–5 years and reaching peak reproductive success  
33 between 6–15 years [Garrott et al., 1991]. On ASIS, however, mares do not normally  
34 breed until they are 3 years old. After that point, fertility rates of 4-year-old mares  
35

37 TABLE 1. Mean ages at death for Assateague horses as a function of length of PZP treatment

	A	B	C	D
Group	Stallions	Untreated mares	Mares treated $< 3$ years	Mares treated $\geq 3$ years
Number	56	42	11	19
Mean age at death (year)	10.33 <sup>b</sup>	6.47 <sup>a</sup>	10.27 <sup>c</sup>	19.94 <sup>c</sup>
SEM	0.845	0.566	1.477	1.666

41 <sup>a</sup>Differs significantly from Groups A,C,D.

43 <sup>b</sup>Differs significantly from Groups B,D.

45 <sup>c</sup>Differs from Groups A–C.



1 were 46%, that of 5-year-old mares were 53%, and that of 6-year-old mares were  
2 69% [Keiper and Houpt, 1984], which is consistent with the data from the study of  
3 Garrott et al. [1991], in western wild horses.

4 The standard treatment protocol in the ASIS management plan applies initial  
5 contraception at 2 years, and booster inoculations at 3 and 4 years of age.  
6 Kirkpatrick and Turner [2002] showed that ASIS mares treated for 3 consecutive  
7 years would take anywhere from 1 to 4+ years to become fertile again, based on  
8 current reversibility data [Kirkpatrick and Turner, 2002], which would cause a  
9 significant decrease in foaling from age 5 to beyond 9 years, which are the years of  
10 peak reproductive success. On ASIS, a study of 14 mares, all 3 years or older, over an  
11 8-year period and before any contraceptive treatment, showed that they produced a  
12 mean of 5.0 foals during that 8-year period [Keiper and Houpt, 1984]. Foals among a  
13 different set of 14 ASIS mares, all 3 years of age or older and treated with PZP for  
14 varying periods of time, produced a mean of 0.5 foals during their lives. Thus, PZP  
15 treatment at 2, 3, and 4 years of age can have a significant negative effect on fertility  
16 and foaling. On ASIS, once a treated animal produces a living foal, she is placed  
17 back on contraception until extinction, further reducing the number of foals  
18 produced in her life.

19 The increased longevity associated with PZP treatment, in PZP-treated  
20 animals, is associated clearly with two phenomena reported earlier [Turner and  
21 Kirkpatrick, 2002], including improved body condition and lower mortality. The  
22 body condition scores of lactating mares have been shown to be significantly lower  
23 than those of non-lactating mares [Rudman and Keiper, 1991]. The  
24 energy requirements for a pregnant or lactating mare are significantly greater than  
25 for open mares. An open mare requires approximately 17.8mcal/day, whereas an  
26 11-month pregnant mare increases to 21.4mcal, and to 30.7mcal in early  
27 lactation [National Research Council, 1989]. These values are for domestic  
28 horses in protected environments and do not take into account the greater  
29 nutritional needs of wild horses living under harsh conditions or game park  
30 animals, where access to food may not be equitable. Thus, it is not surprising  
31 that body condition improves in mares as fertility decreases due to contraception.  
32 The link between increased body condition and increased longevity in horses  
33 is less well understood, but clearly this study suggests some correlation between  
34 the two.

35 In studies of this nature there are several potential sources of error. The first  
36 includes mistakes in the aging of animals, which is usually based on tooth wear. This  
37 approach to aging is particularly tenuous in older age classes. The second source of  
38 error is found in undetected pregnancies because of fetal losses or undetected  
39 neonatal losses. The third source of error involves basing reproductive success on  
40 pregnancy testing or physical condition of mares, to eliminate the undetected  
41 pregnancy and neonatal loss problem. This can include the misclassification of  
42 reproductive condition, based on pregnant mares serum (PMSG) or blood  
43 progesterone analysis for pregnancy detection or inferred lactational condition  
44 based on distended udders.

45 Pregnancy detection using equine chorionic gonadotropin, or PMSG  
46 measurements are relatively accurate through 100 days of pregnancy, but decline  
47 after that. In a study of reproductive rates of western wild horses [Wolfe et al., 1989]  
PMSG analysis was only 53.7% accurate, probably because of the lateness of testing

1 (August through February), well after PMSG levels had declined below the limits of  
the test. The use of serum progesterone analysis alone can also lead to inaccurate  
3 pregnancy rates. In two studies [Wolfe et al., 1989; Garrott et al., 1991] serum  
progesterone analyses resulted in accuracy rates of 85% and 82%, respectively. The  
5 pitfalls using this approach include a failure to distinguish cycling mares from  
pregnant mares, and prolonged elevated progesterone concentrations for weeks to  
7 months after aborted fetuses [Ginther, 1979].

Assessing pregnancy status on the basis of lactating mares or mares with  
9 distended udders has also been shown to be relatively inaccurate. A number of  
investigators have reported the majority of foal mortality during the first 1–2  
11 neonatal months in wild horses [Welsh, 1975; Boyd, 1979; Keiper, 1979], thus mares  
examined after that period would show no distension of the udders and be  
13 misclassified as not having foaled.

None of these potential errors were a factor in the current study, where birth  
15 and death dates were absolutely correct and the presence of foals was confirmed by  
direct observation. Additionally, pregnancy testing on ASIS is based on urinary or  
17 fecal estrone conjugate analysis rather than progesterone or progesterone  
metabolites alone [Kirkpatrick et al., 1986; Daels et al., 1991]. Of 346 horses tested  
19 for pregnancy, 334 proved accurate based on the absence or presence of foals, for a  
96.5% accuracy rate. Of 12 horses whose pregnancy status was not predicted  
21 accurately on the basis of urinary or fecal steroid metabolite analysis, 7 were false  
positives and 5 were false negatives. The 7 false positive horses may in fact have been  
23 accurately pregnant at the time of testing, and experienced fetal loss later in the year.  
Urinary estrone conjugate concentrations, however, decline precipitously within  
25 12 hr of loss of the fetus [Daels et al., 1991]. Additionally, urinary estrone conjugate  
concentrations can distinguish between pregnant animals and cycling animals, and  
27 are useful from approximately Day 40 of pregnancy to parturition [Daels et al., 1991;  
Lasley and Kirkpatrick, 1991]. Previous studies using either urinary estrone  
29 conjugates alone [Kirkpatrick et al., 1990] or with fecal estrone conjugates  
[Kirkpatrick and Turner, 1991a,b; Kirkpatrick et al., 1991] have shown accuracy  
31 approaching 100%. In this retrospective study, pregnancy testing accuracy, over 18  
years, was a minimum of 96.5% accurate. Thus, in this study, error has been  
33 minimized or possibly even eliminated.

## 35 CONCLUSIONS

37 Contraception can be effective in managing free-ranging wildlife or captive  
game park populations [Deigert et al., 2003], and in the process, longevity of  
39 individual animals will be increased significantly, probably across all species. This, in  
turn, will have profound effects on the age profile of the herd, and the intensity of  
41 fertility control application necessary to achieve any given population or growth rate  
goal. At the same time, contraception will improve the health of individual animals  
43 and consequently, the target herd.

## 45 ACKNOWLEDGMENTS

47 Special thanks to J. Kumar, B. Rodgers, G. Olson, and C. Zimmerman, of the  
National Park Service.



## REFERENCES

- Boyd LE. 1979. The mare-foal demography of feral horses in Wyoming's Red Desert. In: Denniston RH, ed. Symposium on the Ecology and Behavior of Wild and Feral Equids, University of Wyoming, Laramie.
- Daels PF, Ammon DC, Stabenfeldt GH, Liu IKM, Hughes JP, Lasley BL. 1991. Urinary and plasma estrogen conjugates, estradiol and estrone concentrations in nonpregnant and early pregnant mares. *Theriogenology* 35:1001-1017.
- Deigert FA, Duncan A, Lyda RO, Frank KM, Kirkpatrick JF. 2003. Immunocontraception of captive exotic species. III. Fallow deer (*Cervus dama*). *Zoo Biol* 22:261-268.
- Dunbar BS, Waldrip NJ, Hedrick J. 1980. Isolation, physicochemical properties and macromolecular composition of zona pellucida from porcine ovaries. *Biochemistry* 19:356-365.
- Fayrer-Husken RA, Grobler D, Van Altena JJ, Kirkpatrick JF, Bertschinger H. 2000. Immunocontraception of free-roaming African elephants. *Nature* 407:149.
- Frank KM, Lyda RO, Kirkpatrick JF. 2005. Immunocontraception of captive exotic species. IV. Species differences in response to the porcine zona pellucida vaccine and the timing of booster inoculations. *Zoo Biol* 24:349-358.
- Garrott RA, Eagle TC, Plotka ED. 1991. Age-specific reproduction in feral horses. *Can J Zool* 69:738-743.
- Ginther OJ. 1979. Reproductive biology of the mare. Cross Plains, WI: Equiserve, Inc. p 334-338.
- Keiper RR. 1979. Population dynamics of feral ponies. In: Denniston RH, ed. Symposium on the Ecology and Behavior of Wild and Feral Equids, University of Wyoming, Laramie.
- Keiper RR, Houtp K. 1984. Reproduction in feral horses: an eight-year study. *Am J Vet Res* 45: 991-995.
- Kirkpatrick JF, Turner JW. 1991a. Changes in herd stallions among feral horse bands and the absence of forced copulation and induced abortion. *Behav Ecol Sociobiol* 29:217-220.
- Kirkpatrick JF, Turner JW. 1991b. Compensatory reproduction among feral horses. *J Wildl Manage* 55:649-652.
- Kirkpatrick JF, Turner A. 2002. Reversibility of action and safety during pregnancy of immunizing against porcine zona pellucida in wild mares (*Equus caballus*). *Reprod Suppl* 60:197-202.
- Kirkpatrick JF, Turner A. 2003. Absence of effects from immunocontraception on seasonal birth patterns and foal survival among barrier island horses. *J Appl Anim Welfare Sci* 6: 301-308.
- Kirkpatrick JF, Liu IKM, Turner JW. 1990a. Remotely-delivered immunocontraception in feral horses. *Wildl Soc Bull* 18:326-330.
- Kirkpatrick JF, Shideler SE, Turner JW. 1990b. Pregnancy determination in uncaptured feral horses based on free steroids in feces and steroid metabolites in urine-soaked snow. *Can J Zool* 68:2576-2579.
- Kirkpatrick JF, Liu IKM, Turner JW, Bernoco M. 1991a. Antigen recognition in mares previously immunized with porcine zona pellucida. *J Reprod Fertil Suppl* 44:321-325.
- Kirkpatrick JF, Shideler SE, Lasley BL, Turner JW. 1991b. Pregnancy determination in uncaptured feral horses by means of fecal steroid conjugates. *Theriogenology* 35:753-759.
- Kirkpatrick JF, Liu IKM, Turner JW, Naugle R, Keiper R. 1992. Long-term effects of porcine zona pellucida immunocontraception on ovarian function of feral horses (*Equus caballus*). *J Reprod Fertil* 93:437-444.
- Kirkpatrick JF, Naugle R, Liu IKM, Bernoco M, Turner JW. 1995a. Effects of seven consecutive years of porcine zona pellucida immunocontraception in feral mares. *Bio Reprod Monogr Series 1: Equine Reproduction* VI. p 411-418.
- Kirkpatrick JF, Zimmermann W, Kolter L, Liu IKM, Turner JW. 1995b. Immunocontraception of captive exotic species: I. Przewalski's horses (*Equus przewalskii*) and banteng (*Bos javanicus*). *Zoo Biol* 14:403-413.
- Kirkpatrick JF, Calle PP, Kalk P, Liu IKM, Bernoco M, Turner JW. 1996. Immunocontraception of captive exotic species. II. Formosan sika deer (*Cervus nippon taiouanus*), Axis deer (*Cervus axis*), Himalayan tahr (*Hemitragus jemlahicus*), Roosevelt elk (*Cervus elaphus roosevelti*), muntjac deer (*Muntiacus reevesi*), and sambar deer (*Cervus unicolor*). *J Zoo Wildl Med* 27: 482-495.
- Lasley BL, Kirkpatrick JF. 1991. Monitoring ovarian function in captive exotic species and free-roaming wildlife by means of urinary and fecal steroids. *J Zoo Wildl Med* 22:23-31.
- National Research Council. 1989. Nutrient requirements of horses. Washington, DC: National Academy of Sciences.
- Naugle RE, Rutberg AT, Underwood HB, Turner JW, Liu IKM. 2002. Field testing of immunocontraception on white-tailed deer (*Odocoileus virginianus*) on Fire Island National Seashore, New York, USA. *Reprod Suppl* 60: 143-153.
- Rudman R, Keiper RR. 1991. The body condition of feral ponies on Assateague Island. *Equine Vet J* 21:455-456.
- Rutberg AT, Naugle RE, Thiele LA, Liu IKM. 2004. Effects of immunocontraception on a suburban population of white-tailed deer *Odocoileus virginianus*. *Biol Conserv* 116: 243-250.
- Shideler SE, Stoops MA, Gee NA, Howell JA, Lasley BL. 2002. Use of porcine zona pellucida (PZP) vaccine as a contraceptive agent in free-ranging mule elk (*Cervus elaphus nansueti*). *Reprod Suppl* 60:169-176.
- Turner JW, Kirkpatrick JF, Liu IKM. 1996. Effectiveness, reversibility and serum antibody

**8 Kirkpatrick and Turner**

- 1     titers associated with immunocastration in  
2     captive white-tailed deer. *J Wildl Manage* 60:  
3     45-51.
- 4     Turner A, Kirkpatrick JF. 2002. Effects of  
5     immunocastration on population, longevity  
6     and body condition in wild mares (*Equus*  
7     *caballus*). *Reprod Suppl* 60:187-195.
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- Welsh AD. 1973. Population, behavioral and  
grazing ecology of the horse of Sable Island,  
Nova Scotia. PhD Thesis, Halifax, Nova Scotia:  
Dalhousie University.
- Wolfe ML, Ellis LC, Macmillan R. 1989. Repro-  
ductive rates of feral horses and burros. *J Wildl*  
*Manage* 53:916-924.